

Driving on sunshine, converting seaweed to furanic biofuels

J.W. van Hal
G. van Hees
mw E.M Cobussen - Pool
Huijgen, W.J.J.

Juni 2017
ECN-L--17-022

Presented at the 6th Congress of the International Society for Applied Phycology,
June 18-23, 2017 in Nantes - France



Driving on sunshine, converting seaweed to furanic biofuels.

Jaap W. van Hal, Guido van Hees,
Esther Cobussen & Wouter J.J.
Huijgen



Acknowledgement



This presentation is part of the MacroFuels project. This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 654010

macrofuels@dti.dk



The Dutch Weed Burger



The Dutch Seaweed Programme

- Seaweed cultivation area 5,000 km² (<10 % of the NL area of the North Sea @ 57,000 km²)
- Integration with off-shore wind parks & (other) aquaculture operations
- Energy potential up to 350 PJ_{th} (25 Mton dry biomass per year)
- Report: ECN-C—05-008



What does ECN do?

- ECN develops market driven technology and know-how to enable a transition to a sustainable energy society
- Business units:
 - Biomass & energy efficiency
 - Solar energy
 - Wind energy
 - Policy studies
 - Environment & energy engineering




ECN

- Independent research institute
- ~500 employees
- Locations:
 - *Petten (HQ)*
 - *Amsterdam*
 - *Eindhoven*
 - *Kuala Lumpur*

What are Furans?


- Class of compounds with a furan-ring.
 - Reaction product of carbohydrate dehydration.
- Generally considered promising biobased building block.
- Challenge:
 - Balance between (acid-catalyzed) furan formation and degradation.



R. Mariscal *et al.* *Energy Environ. Sci.* **2016**, *9* (4), 1144-1189.


Red Macro Algae

Palmaria palmata (Dulse)




About *Palmaria palmata*

- Carbohydrate composition:
 - Rich in xylose, galactose and glucose.
 - Main structural carbohydrate:
 - Xylan polymer (typically ~30wt%).
 - Floridoside (glycerol-galactose heteroside)





Xylan (1,3 and 1,4 linkage)




Floridoside

Forming Furans




Saccharification of *P. palmata*

- Effective saccharification:
- Fresh *P. palmata*
- Catalyst: HCl or commercial xylanase.
- Residual solid: 33-36 dw%.
- Yields monomers using HCl:
 - Xylose up to 85%.
 - Galactose up to 70%.
- Product liquors:
 - Up to 35 g/kg monosaccharides.




Xylose to Furfural

- Single phase (H₂O):
 - Optimisation of process parameters.
 - Brønsted (HCl) and Lewis (SnCl₄) catalysts: at optimum T similar performance.
 - Small positive effect of NaCl on furfural yield.
 - Furfural yield obtained max 60%.
- Biphasic (H₂O/organic):
 - Furfural extracted *in-situ* to prevent degradation.
 - Various extractants tested. Toluene selected for stability and minimal solvent losses.
 - Furfural yield increases to near theoretical (HCl).



In one step!

- Single step:
 - Water:
 - Furfural yield 38% (0.2M HCl, 1h, 170 °C).
 - Water-toluene:
 - Furfural yield 75% (0.3M HCl / 0.9M NaCl, 1h, 170 °C, 10wt% *P. palmata*).
- Two steps:
 - Hydrolysis of seaweed polysaccharides to monomers.
 - Dehydration of xylose to furfural in hydrolysate.
 - Biphasic process hydrolysate/toluene 1:2 v/v.
 - No additional acid used.
 - Overall yield from *P. palmata* to furfural: 98%.
 - No negative matrix effects observed.




Green Macroalgae

Ulva sp.



About *Ulva lactuca*


- Why *Ulva*?
 - Unique carbohydrate composition, incl. rhamnose.
 - Ulvan (rhamnose, xylose, glucuronic acid, iduronic acid).
 - Cellulose (glucose).
 - Dehydration of rhamnose yields 5-methylfurfural.
 - Directly applicable as biofuel (additive).



Forming 5-methyl furfural




Ulva lactuca




Saccharification *Ulva lactuca*

- Hydrolysis of polysaccharides to monomeric carbohydrates demonstrated with fresh seaweed.
- Monomeric yields of major carbohydrates (Glucose, Rhamnose, and Xylose) of at least 85% possible.
- However, low sugar concentrations in product liquors (~5 g/kg) due to low carbohydrate content seaweed.



Rhamnose to 5-methylfurfural

- Scant information dehydration of rhamnose in the literature.
- Similar approach and conditions applied as for *P. palmata*.
- Direct HCl-catalyzed dehydration in water:
- Low yield of 5-methylfurfural (max 22%).




1h, 0.3M HCl, 0.5M NaCl, water/toluene

U. lactuca to 5-methylfurfural

- Conversion of *U. lactuca* more challenging than *P. palmata*:
 - Poor 5-methylfurfural yield achieved directly in water: 25%.
 - Biphasic system with toluene: 36%.
 - Two-step approach (saccharification & dehydration): 56%.
- Simultaneous conversion of other ulvan building blocks (such as xylose).

In pictures



Conclusions

- Effective saccharification of *P. palmata* and *U. lactuca* feasible.
- Effective conversion of seaweed carbohydrates to furans feasible when applying in-situ extraction.
- *P. palmata* most suited seaweed for carbohydrate or furan production.
 - Higher carbohydrate content.
 - Furfural yields higher than 5-methylfurfural yields.

In numbers

Process / yields	<i>P. palmata:</i> Xyl → furfural	<i>U. lactuca:</i> Rham → 5-methylfurfural
One-step approach in H ₂ O	38	25
One-step approach in H ₂ O/toluene	75	36
Two-step approach with H ₂ O/toluene	98	56

Thank for your attention!

Questions?

Publications:

<https://www.ecn.nl/publications/>

<http://www.macrofuels.eu>

<http://www.macrocascade.eu>

<http://www.noordzeeboerderij.nl>



Jaap W. van Hal, Ph.D.

Innovation Manager Biorefinery

T +31 88 515 4297

M +31 6 25382913

vanhal@ecn.nl

P.O. Box 1, 1755 ZG PETTEN

The Netherlands

www.ecn.nl

ECN

Westerduinweg 3
1755 LE Petten
The Netherlands

P.O. Box 1
1755 ZG Petten
The Netherlands

T +31 88 515 4949
F +31 88 515 8338
info@ecn.nl
www.ecn.nl

